

OTR FILE COPY

NPS-53-88-009

NAVAL POSTGRADUATE SCHOOL

Monterey, California

AD-A199 345



DTIC
ELECTE
OCT 1 2 1988
S H D

A NEW CONSERVATIVE SCHEME
FOR SOLVING THE TWO BODY PROBLEM

B. Neta
Y. Ilan-Lipowski

August 1988

Approved for public release; distribution unlimited
Prepared for:

Naval Postgraduate School
Monterey, CA 93943

88 1012 022


MONTEREY, CALIFORNIA

Rear Admiral R. C. Austin
Superintendent

Harrison Shull
Provost


Reproduction of all or part of this report is authorized.


This report was prepared by:


BENY NETA
Department of Mathematics

Reviewed by:

Released by:


HAROLD M. FREDRICKSEN
Chairman
Department of Mathematics


KNEALE T. MARSHALL
Dean of Information and
Policy Sciences

REPORT DOCUMENTATION PAGE

13 REPORT SECURITY CLASSIFICATION UNCLASSIFIED			10 RESTRICTIVE MARKINGS			
2a SECURITY CLASSIFICATION AUTHORITY			3 DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited			
2b DECLASSIFICATION/DOWNGRADING SCHEDULE						
4 PERFORMING ORGANIZATION REPORT NUMBER(S) NPS-53-88-009			5 MONITORING ORGANIZATION REPORT NUMBER(S) NPS-53-88-009			
6a NAME OF PERFORMING ORGANIZATION Naval Postgraduate School		6b OFFICE SYMBOL (if applicable) 53		7a NAME OF MONITORING ORGANIZATION		
6c ADDRESS (City, State, and ZIP Code) Monterey, CA 93943				7b ADDRESS (City, State, and ZIP Code)		
8a NAME OF FUNDING/SPONSORING ORGANIZATION NSWC		8b OFFICE SYMBOL (if applicable) K-13		9 PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER		
6c ADDRESS (City, State, and ZIP Code) Dahlgren, VA 22448				10 SOURCE OF FUNDING NUMBERS		
				PROGRAM ELEMENT NO	PROJECT NO	TASK NO
11 TITLE (Include Security Classification) A New Conservative Scheme for Solving the Two Body Problem						
12 PERSONAL AUTHOR(S) B. Neta and Y. Ilan-Lipowski						
13a TYPE OF REPORT Technical Report		13b TIME COVERED FROM 10/86 TO 6/87		14 DATE OF REPORT (Year Month Day) August 1988		15 PAGE COUNT
16 SUPPLEMENTARY NOTATION						
17 COSATI CODES			18 SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Two Body Problem, Conservative Scheme			
FIELD	GROUP	SUB-GROUP				
19 ABSTRACT (Continue on reverse if necessary and identify by block number) The well-known Stormer-Cowell class of linear k-step methods for the solution of second order initial value problems suffer from orbital instability. The solution of a problem describing a uniform motion in a circular orbit, spirals inward. Several modifications were suggested, some require the a-priori knowledge of the frequency. Here we develop a method based on the conservation of energy and momentum. This method overcomes the aforementioned difficulty.						
20 DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS				21 ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a NAME OF RESPONSIBLE INDIVIDUAL B. Neta & Y. Ilan-Lipowski				22b TELEPHONE (include Area Code) 408-646-2234		22c OFFICE SYMBOL 53Nd

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted
All other editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE
UNCLASSIFIED

A New Conservative Scheme
for Solving the Two Body Problem

B. Neta
Y. Ilan-Lipowski

Naval Postgraduate School
Monterey, CA 93943

ACKNOWLEDGEMENT

The authors would like to thank NSWC for its partial support of this research.

Abstract

The well-known Störmer-Cowell class of linear k-step methods for the solution of second order initial value problems suffer from orbital instability. The solution of a problem describing a uniform motion in a circular orbit, spirals inward. Several modifications were suggested, some require the a-priori knowledge of the frequency. Here we develop a method based on the conservation of energy and momentum. This method overcomes the aforementioned difficulty.

1980 Mathematics Subject Classification 65L05

Key Words: Conservative scheme, Two body problem.



Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	<input type="checkbox"/>
Distribution	
Availability	
Date	
A-1	

1. Introduction

In this paper we develop a method for the numerical solution of the equations of motion of an object acted upon by several gravitational masses. In general, the motion can be described by a special class of second order differential equations, namely

$$y''(x) = f(x, y(x)) . \quad (1)$$

There exist methods of Runge-Kutta type which tackle this problem directly (Collatz [8, p. 61], de Vogelaere [23], Scraton [20] and others). Also, linear multistep methods of the form

$$\sum_{j=0}^k \alpha_j y_{n+j} = h^2 \sum_{j=0}^{k'} \beta_j f_{n+j} \quad (2)$$

exist. See, for example, Henrici [11, p. 289], Lambert [13, p. 252] and others. One of the authors has developed hybrid methods for such classes (Neta and Lee [15], Neta [16, 18]).

The direct application of methods of class (2) to problem (1), rather than the application of a conventional linear multistep method to an equivalent first-order system is usually recommended (Ash [1]).

Special methods based on a-priori knowledge of the frequency were developed by Bettis [3], Steifel and Bettis [22], Gautschi [10], Neta and Ford [17], Neta [19], van der Houwen and Sommeijer [12], Lyche [14] and Sommeijer et al [21].

To avoid the use of the frequency other methods were developed satisfying the so-called P-stability. This idea of P-stability was introduced by Lambert and Watson [13]. They showed that such methods are necessarily implicit and cannot have order greater than two. Chawla [5] and Cash [4] independently showed that this order-barrier can be crossed over by considering hybrid two-step methods. Other P-stable methods were developed by Costabile & Costabile [9], Chawla [6], Chawla and Rao [7] and others.

Our idea here is to develop a new method to approximate the solution of the two body problem. This scheme will conserve both the energy per unit mass and the specific angular momentum of the system.

In the next section the method is developed for the perturbation free flight. Numerical experiments with the scheme will be described in the last section.

2. Development of the Method

In this section we describe our new scheme to solve the following system of equations

$$\frac{d^2x}{dt^2} = - \frac{k}{(x^2+y^2)^{3/2}} x, \quad (3)$$

$$\frac{d^2y}{dt^2} = - \frac{k}{(x^2+y^2)^{3/2}} y, \quad (4)$$

where $k = 3.9877 \cdot 10^{14} \text{ m}^3 \text{ sec}^{-2}$ is the gravitational parameter.

It is well known that Cowell's method gives a numerical solution that spirals inward. Encke's method improves this result but requires more work [2].

It can be easily shown (see e.g., Bate et al. [2]) that the energy and momentum are conserved for a perturbation free flight. The conservation of these quantities will be used to obtain the fast changing variable. It is thus important to rewrite the system in polar coordinates

$$\frac{d^2r}{dt^2} = r \left(\frac{d\theta}{dt} \right)^2 - \frac{k}{r^2} \quad (5)$$

$$\frac{d^2\theta}{dt^2} = - \frac{2}{r} \frac{dr}{dt} \frac{d\theta}{dt}. \quad (6)$$

The radius r does not change much in time and thus we can use any method; here we use a second order Taylor series method. The

value of θ corresponding to r will be obtained by integrating the relation (conservation of energy per unit mass)

$$\left(\frac{dr}{dt}\right)^2 + \left(r \frac{d\theta}{dt}\right)^2 = \left(\frac{dr}{dt}\right)_{t=0}^2 + (r(0) \frac{d\theta}{dt})_{t=0}^2. \quad (7)$$

To be more specific, our method approximates r using

$$r(t_{i+1}) = r(t_i) + \dot{r}(t_i)(\Delta t) + \frac{1}{2}\ddot{r}(t_i)(\Delta t)^2 \quad (8)$$

where

$$\dot{r}(t_{i+1}) = \dot{r}(t_i) + \ddot{r}(t_i)(\Delta t) \quad (9)$$

and $\ddot{r}(t)$ is obtained from the differential equation. In a similar fashion we find

$$\theta(t_{i+1}) = \theta(t_i) + \dot{\theta}(t_i)(\Delta t) + \frac{1}{2}\ddot{\theta}(t_i)(\Delta t)^2 \quad (10)$$

where

$$\dot{\theta}(t_i) = \sqrt{(C + k/r(t_i))^2 - \dot{r}^2(t_i)/r(t_i)} \quad (11)$$

and $\ddot{\theta}(t_i)$ is obtained by differentiating the above relation.

The results of this method were compared to Taylor series method for both r and θ where now $\dot{\theta}$ is evaluated from an equation similar to (9).

Remark: It should be clear that this idea can be used for any conservative system. One uses the conservation of some quantity to obtain an approximate value for the faster changing component.

3. Numerical Experiments

In this section we solve the equations of motion (3)-(4) combined with the initial values

$$\begin{aligned} r(0) &= R + 10^5, \\ r'(0) &= 0, \\ \epsilon(0) &= 0, \\ \dot{\epsilon}(0) &= 8000/r(0), \end{aligned} \tag{12}$$

where $R = 6.371 \times 10^6$ is the radius of earth.

In the first table we compare the results of our method with Taylor series using $t = 1$ sec. As can be seen in the table, the value of r at the end of each period is nearly constant with our method but decreases as the number of periods increases with Taylor series method.

TABLE 1

At the end of Period	$r \cdot 10^7$ (10 ⁷ m)		Energy per unit mass (10 ⁸ m ² /sec ²)	Specific angular momentum (10 ¹¹ m ² /sec)
	Taylor	New		
0	.6471	.6471	-.2962	.5277
5	.6320	.6476	-.3019	.5127
10	.6173	.6482	-.3077	.5078
15	.6030	.6487	-.3135	.5029
20	.5890	.6493	-.3194	.4982
30	.5622	.6503	-.3313	.4888
40	.5367	.6513	-.3433	.4798
50	.5123	.6523	-.3557	.4710
60	.4891	.6533	-.3683	.4624
70	.4667	.6543	-.3811	.4540

The energy per unit mass E and the specific angular momentum are given by

$$E = \frac{1}{2}(\dot{r}^2 + (r\dot{\theta})^2) - \frac{k}{r} , \quad (13)$$


$$P = r^2\dot{\theta} - r\dot{\theta} .$$

In our example both the energy and momentum are conserved.


In the following three figures the radius r , the energy per unit mass E and the specific angular momentum P were plotted at the end of every 10 periods. It is clear that the relative change in r values after 70 periods using Taylor series method is more than 25 times larger compared to our new method. The energy and momentum were kept constant by the new scheme whereas the Taylor series method shows a relative loss of 28% in energy and 12% in momentum at the end of 70 periods.

Remark: Note that our method is explicit whereas the P-stable methods suggested by Lambert and Watson [13] are implicit.


COMPARISON OF RADIUS



COMPARISON OF ENERGY



COMPARISON OF MOMENTUM



References

1. J.H. Ash, Analysis of multistep methods for second-order ordinary differential equations, Ph.D. thesis, University of Toronto (1969).
2. R.R. Bate, D.D. Mueller, J.E. White, Fundamentals of Astrodynamics, Dover Publications, New York, 1971.
3. D.G. Bettis, Numerical integration of product of Fourier and ordinary polynomials, Numer. Math., 14, 421-434 (1970).
4. J.R. Cash, High-order P-stable formulae for the numerical integration of periodic initial value problems, Numer. Math., 37, 355-370 (1981).
5. M.M. Chawla, Two step fourth-order P-stable methods for second order differential equations, BIT, 21, 190-193 (1981).
6. M.M. Chawla, Unconditionally stable Nourmerov-type methods for second order differential equations, BIT, 23, 541-542 (1983).
7. M.M. Chawla, P.S. Rao, High-accuracy P-stable methods for $y'' = f(t, y)$, IMA J. Numer. Anal., 5, 215-220 (1985).
8. L. Collatz, Numerical Treatment of Differential Equations, Springer, 1960.
9. F. Costabile, C. Costabile, Two-step fourth order P-stable methods for second order differential equations, BIT, 22, 384-386 (1982).
10. W. Gautschi, Numerical integration of ordinary differential equations based on trigonometric polynomials, Numer. Math., 3, 381-397 (1961).
11. P. Henrici, Discrete Variable Methods in Ordinary Differential Equations, Wiley, 1962.
12. P.J. van der Houwen, B.P. Sommeijer, Linear multistep methods with minimized truncation error for periodic initial value problems, IMA J. Numer. Anal., 4, 479-489 (1984).
13. J.D. Lambert, I.A. Watson, Symmetric multistep methods for periodic initial value problems, J. Inst. Maths Applics, 18, 189-202 (1976).
14. T. Lyche, Chebyshevian multistep methods for ordinary differential equations, Numer. Math., 19, 65-72 (1972).

15. B. Neta, S.C. Lee, Hybrid methods for a special class of second-order differential equations, Cong. Numer., 38, 203-225 (1983).
16. B. Neta, Higher order hybrid Stormer-Cowell methods for ordinary differential equations, Cong. Numer., 42, 251-264 (1984).
17. B. Neta, C.H. Ford, Families of methods for ordinary differential equations based on trigonometric polynomials, J. Comp. Appl. Math., 10, 33-38 (1984).
18. B. Neta, Hybrid predictors and correctors for solving a special class of second order differential equations, Cong. Numer., 46, 241-248 (1985).
19. B. Neta, Families of backward differentiation methods based on trigonometric polynomials, to appear.
20. R.E. Scraton, The numerical solution of second-order differential equations not containing the first derivative explicitly, Comput. J., 6, 368-370 (1964).
21. B.P. Sommijs, P.J. van der Houwen, B. Neta, Symmetric linear multistep methods for second-order differential equations with periodic solutions, Appl. Numer. Math., 2, 69-77 (1986).
22. E. Steifel, D.G. Bettis, Stabilization of Cowell's method, Numer. Math., 13, 154-175 (1969).
23. R. de Vogelaere, A method for the numerical integration of differential equations of second-order without explicit first derivatives, J. Res. Nat. Bur. Standards, 54, 119-125 (1955).

INITIAL DISTRIBUTION LIST

DEFENSE TECHNICAL INFO CTR (2)
CAMERON STATION
ALEXANDRIA, VA 22304-6145

PROFESSOR M. F. PLATZER
NAVAL POSTGRAD. SCHOOL
DEPT OF AERONAUTICS
CODE 67P1
MONTEREY, CA 93943

Y. ILAN-LIPOWSKI (5)
27 ZIMCHE HAYEHUDIM STREET
TEL AVIV, ISRAEL

DIRECTOR (2)
DEFENSE TECH INFO CTR,
CAMERON STATION
ALEXANDRIA, VA 22314

LIBRARY (2)
NAVAL POSTGRAD. SCHOOL
MONTEREY, CA 93943

CENTER FOR NAVAL ANALYSES
4401 FORD AVENUE
ALEXANDRIA, VA 22302-0268

PROFESSOR B. NETA (15)
NAVAL POSTGRAD. SCHOOL
DEPT OF MATHEMATICS
CODE 53Nd
MONTEREY, CA 93943

PROFESSOR M. WEIR
NAVAL POSTGRAD. SCHOOL
DEPT OF MATHEMATICS
CODE 53Wc
MONTEREY, CA 93943

DR. DAN PARKS
NAVAL SURF. WEAPONS CTR
K-13
DAHLGREN, VA 22448

DIR. OF RESEARCH ADMIN
CODE 012
NAVAL POSTGRAD. SCHOOL
MONTEREY, CA 93943

DEPT OF MATHEMATICS
NAVAL POSTGRAD. SCHOOL
MONTEREY, CA 93943